

FLUVAC INNOVATOR[®] 4

Fort Dodge

Encephalomyelitis-Influenza Vaccine

Eastern & Western, Killed Virus

Tetanus Toxoid

INDICATIONS: For intramuscular vaccination of healthy horses as an aid in the prevention of equine encephalomyelitis due to Eastern and Western viruses, equine influenza due to type A₂ viruses, and tetanus.

DOSAGE AND ADMINISTRATION: Horses, inject one 1 mL dose intramuscularly using aseptic technique. Administer a second 1 mL dose 3 to 4 weeks after the first dose. A 1 mL booster dose should be given annually. Early revaccination may be advisable when horses are faced with an outbreak or with other conditions which might make heavy exposure likely.

CAUTION: Store in the dark at 2° to 7°C (35° to 45°F). AVOID FREEZING. SHAKE WELL. Use entire contents when first opened. In some instances, transient local reactions may occur at the injection site. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction, administer epinephrine.

Thimerosal, neomycin and polymyxin B added as preservatives.

For Veterinary Use Only

Remove serial number and expiration date label and place on the animal's medical chart.

© 2004 Fort Dodge Animal Health. All Rights Reserved.

Fort Dodge Animal Health, Fort Dodge, Iowa 50501 U.S.A.

U.S. Vet. License No. 112

| | |
|------------------------------|--------|
| 12 x 1 mL preloaded syringes | 18061C |
| 10 x 10 mL (10 dose) vials | 18063B |

NAC No.: 10032764